



**NANO-HYDROLYZED  
COLLAGEN PROTEIN**



Serving size: 1 ounce/ 30 mL or 1 individual packet

**Administer orally or through feeding tube**

<b>Indications</b>	
Pressure Injuries (all stages)	Post-Surgery
Wounds (diabetic, surgical, burns)	Fluid Restriction
Protein Energy Malnutrition	Clear Liquid Diet
Inadequate Protein Intake	Low Serum Proteins
Dialysis	Bariatric Surgery

**ProT Gold is great without mixing or diluting but may be added to beverages or food as desired.**

<b>Easily Mix with:</b>	
Water	Oatmeal/ Hot Cereal
Juice	Yogurt
Sparkling Water/ Soda	Applesauce/ Fruit Puree
Hot or Cold Beverage of Choice	Smoothies/ Shakes

**See Reverse for Tube Feeding Procedures**



# ENTERAL ADMINISTRATION PROCEDURE

**Do not add ProT Gold to an open or closed tube feeding formula system or directly into enteral formula.**

**Do not add medications to ProT Gold.**

<b>1</b>	Flush feeding tube with 30-60 mL of water	
<b>2</b>	<b><u>30 oz Bottle:</u></b> Measure proper dosage using 1-2 oz. dosing cup or 3-6 oz. container if dilution is needed.	<b><u>1 oz. Packet:</u></b> Empty entire contents of package into enteral syringe with feeding tube clamped or 3-6 oz. container if dilution is needed.
<b>3</b>	It may be necessary to dilute ProT Gold with 15-60 mL of water for smaller tube sizes.	
<b>4</b>	Administer ProT Gold using enteral syringe directly into feeding tube.	
<b>5</b>	Flush feeding tube with 30-60 mL of water.	

**Store at room temperature.**

**Record date on bottom of container upon opening 30 oz. bottles. Discard opened container after 90 days.**

**ProT Gold is not suitable for sole source nutrition and is not for parenteral use.**